

Formelsammlung

Orthogonalpolynome

a	b	$\omega(x)$	Name	Formel
-1	1	$(1-x)^\alpha(1+x)^\beta$	Jacobi-P.	$P_n^{(\alpha,\beta)}(x) = \frac{1}{2^n n! \omega(x)} \frac{d^n}{dx^n} (\omega(x)(x^2-1)^n)$
0	∞	e^{-x}	Laguerre-P.	$\ell_n(x) = \frac{e^x}{n!} \frac{d^n}{dx^n} (e^{-x} x^n)$
$-\infty$	∞	e^{-x^2}	Hermite-P.	$H_n(x) = e^{x^2} \frac{d^n}{dx^n} (e^{-x^2})$

Eigenschaften der Jacobi-Polynome

Orthogonalität: (mit der Γ -Funktion)

$$\int_{-1}^1 (1-x)^\alpha(1+x)^\beta P_n^{(\alpha,\beta)}(x) P_m^{(\alpha,\beta)}(x) dx = \delta_{mn} \frac{2^{\alpha+\beta+1} \Gamma(n+\alpha+1) \Gamma(n+\beta+1)}{(2n+\alpha+\beta+1) n! \Gamma(n+\alpha+\beta+1)}$$

3-Term-Rekursion:

$$2(n+1)(n+\alpha+\beta+1)(2n+\alpha+\beta) P_{n+1}^{(\alpha,\beta)}(x) = (2n+\alpha+\beta+1)[(\alpha^2-\beta^2) + x(2n+\alpha+\beta)(2n+\alpha+\beta+2)] P_n^{(\alpha,\beta)}(x) - 2(n+\alpha)(n+\beta)(2n+\alpha+\beta+2) P_{n-1}^{(\alpha,\beta)}(x)$$

Nullstellen: Eigenwerte der tridiagonalen Matrix

$$\mathcal{T}_n = \begin{bmatrix} a_0 & b_1 & 0 & \dots & 0 \\ b_1 & a_1 & b_2 & 0 & \dots \\ 0 & b_2 & a_2 & b_3 & \\ \vdots & & & \ddots & b_{n-1} \\ 0 & \dots & 0 & b_{n-1} & a_{n-1} \end{bmatrix} \in \mathbb{R}^{n,n}$$

mit

$$a_j = \frac{(\beta-\alpha)(\beta+\alpha)}{(2j+\alpha+\beta)(2j+2+\alpha+\beta)},$$

$$b_j = \sqrt{\frac{4j(j+\alpha)(j+\beta)(j+\alpha+\beta)}{(2j-1+\alpha+\beta)(2j+\alpha+\beta)^2(2j+1+\alpha+\beta)}}$$

Speziell ist $P_n^{(0,0)}(x) = L_n(x)$ (Legendre-Polynom) und $P_n^{-1/2,-1/2}(x) = T_n(x)$ (Tschebyscheffpolynom).

Moore-Penrose-Inverse

Die Moore-Penrose Inverse A^\dagger von $A \in \mathbb{C}^{m \times n}$ ist die eindeutige Lösung $X \in \mathbb{C}^{n \times m}$ der 4 Gleichungen

$$\begin{aligned} AXA &= A, \\ XAX &= X, \\ (AX)^* &= AX, \\ (XA)^* &= XA. \end{aligned}$$

Orthogonale Transformationen

Householder

Es sei $a = [a_i]_{i=1}^n$. Mit

$$v = \begin{cases} \frac{1}{|a_1| \|a\|_2} (|a_1| a + a_1 \|a\|_2 e_1) & a_1 \neq 0 \\ a / \|a\|_2 + e_1 & a_1 = 0 \end{cases}$$

wird $Pa = \beta e_1$ für $P = I - 2 \frac{vv^*}{v^*v}$.

Givens-Rotation

Mit

$$t = \frac{x_2}{|x_1|}, n = \sqrt{1 + |t|^2}, \quad c = \frac{x_1/|x_1|}{n}, s = \frac{t}{n} \quad |x_1| \geq |x_2|,$$

$$t = \frac{x_1}{|x_2|}, n = \sqrt{1 + |t|^2}, \quad s = \frac{x_2/|x_2|}{n}, c = \frac{t}{n} \quad |x_1| < |x_2|,$$

setzen wir

$$C_{cs} = C_{\theta, \alpha, \beta} = \begin{bmatrix} \bar{c} & \bar{s} \\ -s & c \end{bmatrix} \quad c = e^{i\alpha} \cos \theta, s = e^{i\beta} \sin \theta, \quad \alpha, \beta \in [0, 2\pi).$$

Dann ist

$$C_{cs}x = \|x\|_2 e_1.$$

Hutfunktionenbasis auf $[0, 1]$

Auf Netz mit den Knoten

$$x_i = ih, \quad h = \frac{1}{n}, i = 0, \dots, n$$

sind

$$\phi_i(x) = \begin{cases} n(x - x_{i-1}) & x \in [x_{i-1}, x_i] \\ n(x_{i+1} - x) & x \in [x_i, x_{i+1}] \\ 0 & \text{sonst} \end{cases} \quad i = 1, \dots, n-1,$$

PCG-Verfahren

$$x^{(k+1)} = x^{(k)} + \alpha_k q^{(k)}, \quad \text{mit } \alpha_k = \frac{\langle w^{(k)}, r^{(k)} \rangle}{\langle q^{(k)}, Aq^{(k)} \rangle}, \quad k \geq 0$$

$$r^{(k+1)} = \begin{cases} r^{(k)} + \alpha_k Aq^{(k)} & k \geq 0 \\ Ax^{(0)} - b & k = -1 \end{cases},$$

$$w^{(k+1)} = C^{-1}r^{(k+1)}, \quad k \geq 0$$

$$q^{(k+1)} = \begin{cases} \beta_k q^{(k)} - w^{(k+1)} & k \geq 0 \\ w^{(0)} & k = -1 \end{cases}, \quad \text{mit } \beta_k = \frac{\langle w^{(k+1)}, r^{(k+1)} \rangle}{\langle w^{(k)}, r^{(k)} \rangle}$$

Arnoldi-Prozeß in Krylowraum

$$v_{j+1} = Aq_j - \sum_{i=1}^j (q_i^* Aq_j) q_i$$

$$q_{j+1} = \|v_{j+1}\|_2^{-1} v_{j+1}$$

Als Matrixschreibweise:

$$AQ_{k-1} = Q_k H_{k-1}$$

mit der oberen Hessenbergmatrix

$$H = \begin{bmatrix} h_{11} & h_{12} & h_{13} & \dots & h_{1,k-1} \\ h_{21} & h_{22} & h_{23} & \dots & h_{2,k-1} \\ 0 & h_{32} & h_{33} & \dots & h_{3,k-1} \\ \dots & \vdots & \vdots & & \\ 0 & 0 & \dots & 0 & h_{k,k-1} \end{bmatrix} \in \mathbb{C}^{k \times (k-1)}, \quad h_{ij} = \begin{cases} q_i^* Aq_j & i \leq j \\ v_i^* v_i & i = j+1 \\ 0 & \text{sonst} \end{cases}$$

Gauß-Legendre-Formeln mit n Stützstellen und Gewichten (x_i, ω_i) , $i = 0, \dots, n-1$

Anzahl Punkte	Stützstellen	Gewichte
1	$x_0 = 0$	$\omega_0 = 2$
2	$x_{0/1} = \pm \sqrt{1/3}$	$\omega_{0/1} = 1$
3	$x_{0/1} = \pm \sqrt{3/5}, x_2 = 0$	$\omega_{0/1} = \frac{5}{9}, \omega_2 = \frac{8}{9}$
4	$x_{0/1} = \frac{1}{7} \pm \sqrt{3 - 2\sqrt{6/5}}, x_{2/3} = \frac{1}{7} \pm \sqrt{3 + 2\sqrt{6/5}}$	$\omega_{0/1} = \frac{18 + \sqrt{30}}{36}, \omega_{2/3} = \frac{18 - \sqrt{30}}{36}$
5	$x_{0/1} = \frac{1}{3} \pm \sqrt{5 - 2\sqrt{10/7}}, x_{2/3} = \frac{1}{3} \pm \sqrt{5 + 2\sqrt{10/7}}, x_4 = 0$	$\omega_{0/1} = \frac{322 + 13\sqrt{70}}{900}, \omega_{2/3} = \frac{322 - 13\sqrt{70}}{900}, \omega_4 = \frac{128}{225}$