



Institut für Numerische Simulation

Rheinische Friedrich-Wilhelms-Universität Bonn

Wegelerstraße 6 • 53115 Bonn • Germany
phone +49 228 73-3427 • fax +49 228 73-7527
www.ins.uni-bonn.de

M. Griebel, H. Harbrecht

**Approximation of two-variate functions:
Singular value decomposition versus sparse grids**

INS Preprint No. 1109

2011

Approximation of two-variate functions: singular value decomposition versus sparse grids

MICHAEL GRIEBEL[†]

Institut für Numerische Simulation, Universität Bonn, Wegelerstr. 6, 53115 Bonn, Germany

AND

HELMUT HARBRECHT[‡]

Mathematisches Institut, Universität Basel, Rheinsprung 21, 4051 Basel, Switzerland

[Received on 6 November 2012]

We compare the cost complexities of two approximation schemes for functions $f \in H^p(\Omega_1 \times \Omega_2)$ which live on the product domain $\Omega_1 \times \Omega_2$ of sufficiently smooth domains $\Omega_1 \subset \mathbb{R}^{n_1}$ and $\Omega_2 \subset \mathbb{R}^{n_2}$, namely the singular value / Karhunen-Löve decomposition and the sparse grid representation. Here we assume that suitable finite element methods with associated *fixed* order r of accuracy are given on the domains Ω_1 and Ω_2 . Then, the sparse grid approximation essentially needs only $\mathcal{O}(\varepsilon^{-q})$ with $q = \frac{\max\{n_1, n_2\}}{r}$ unknowns to reach a prescribed accuracy ε provided that the smoothness of f satisfies $p \geq r \frac{n_1 + n_2}{\max\{n_1, n_2\}}$, which is an almost optimal rate. The singular value decomposition produces this rate only if f is analytical since otherwise the decay of the singular values is not fast enough. If $p < r \frac{n_1 + n_2}{\max\{n_1, n_2\}}$, then the sparse grid approach gives essentially the rate $\mathcal{O}(\varepsilon^{-q})$ with $q = \frac{n_1 + n_2}{p}$ while, for the singular value decomposition, we can only prove the rate $\mathcal{O}(\varepsilon^{-q})$ with $q = \frac{2 \min\{r, p\} \min\{n_1, n_2\} + 2p \max\{n_1, n_2\}}{(2p - \min\{n_1, n_2\}) \min\{r, p\}}$. We derive the resulting complexities, compare the two approaches and present numerical results which demonstrate that these rates are also achieved in numerical practice.

Keywords: singular value decomposition; sparse grids; complexity.

1. Introduction

The efficient approximate representation of multivariate functions is an important task in numerical analysis and scientific computing. In this paper, we concentrate on functions which live on the product of two domains $\Omega_1 \times \Omega_2$. Already for this simple situation, there exists a large amount of applications. For example, radiosity models and radiative transfer (Widmer *et al.* (2008)) exhibit a product structure. Here, Ω_1 denotes the spatial three-dimensional domain of the geometric object under consideration and Ω_2 is the sphere \mathbb{S}^2 . Moreover, in the case of space-time discretizations of parabolic problems, Ω_1 is the time interval whereas Ω_2 is the spatial domain (Griebel & Oeltz (2007) and Stevenson & Schwab (2009)).

Then, there are various phase space problems where both Ω_1 and Ω_2 are three-dimensional cubes or the full three-dimensional real space. Examples are the Boltzmann equation, kinetic equations or the Langevin equation, see e.g. Balescu (1997). Furthermore, non-Newtonian flow can be modeled by a coupled system which consists of the Navier Stokes equation for the flow in a three-dimensional

[†]griebel@ins.uni-bonn.de

[‡]helmut.harbrecht@unibas.ch

geometry described by Ω_1 and of the Fokker-Planck equation in a $3(k-1)$ -dimensional configuration space on Ω_2 . Here k denotes the number of atoms in a chain-like molecule which constitutes the non-Newtonian behavior of the flow, for details see Barrett *et al.* (2009), Le Bris & Lelièvre (2009), and Lozinski *et al.* (2010). Note that the domain of the configuration space is itself again a product of $k-1$ spheres.

Another example is two-scale homogenization. After unfolding (Cioranescu *et al.* (2008)), it gives raise to the product of the macroscopic physical domain and the periodic microscopic domain of the cell problem, see Matache (2002). For multiple scales, a general product appears here which still can be written as the product of two domains, one containing e.g. the macroscopic scale and the other consisting of the product of the domains of the different microscales (Hoang & Schwab (2005)).

Also the two-point correlation functions of linear elliptic boundary value problems with stochastic source terms, i.e. $Au(\omega) = f(\omega)$ in Ω , are known to satisfy a deterministic partial differential equation on the product domain $\Omega \times \Omega$. Namely, the two-point correlation satisfies the equation $(A \otimes A) \text{Cor}_u = \text{Cor}_f$ in $\Omega \times \Omega$, see Schwab & Todor (2003a). Higher order moments then involve larger tensor products (Schwab & Todor (2003b)). This approach extends to stochastic diffusion functions and more general PDEs with stochastic coefficient functions as well as to stochastic domains (Harbrecht *et al.* (2008a), Harbrecht (2010)).

Finally we find the product of two domains in quantum mechanics for e.g. the Schrödinger equation for Helium; systems with more than two electrons involve then multiple product domains, of course.

In general, some problems are directly given on the product of two domains, while for other problems the domains themselves are products of lower-dimensional domains. Then, the domain of an n -dimensional problem with e.g. n being some power of two can be split into the tensor product of two domains of dimension $n/2$ which can be recursively further split until a terminal situation (a one-dimensional domain or a truly higher dimensional but non-tensor product domain) is reached. Related representation methods have recently been considered in Beberdorf (2011), Grasedyck (2010), Hackbusch (2012), Hackbusch & Kühn (2009), or Oseledets & Tyrtshnikov (2009).

In this article, we consider the simple case of two domains Ω_1 and Ω_2 only. Here, our analysis covers the situation of the first bisection step in the above mentioned recursion. To this end, for $i = 1, 2$, let Ω_i denote a domain in \mathbb{R}^{n_i} (or alternatively also an n_i -dimensional manifold in \mathbb{R}^{n_i+1}). We consider the numerical approximation of two-variate functions. That is, on the individual subdomains we suppose to have given, fixed sequences of nested trial spaces

$$V_0^{(i)} \subset V_1^{(i)} \subset V_2^{(i)} \subset \dots \subset L^2(\Omega_i)$$

which consist of ansatz functions of the same, fixed approximation order r . Based on these ansatz spaces, we intend to compare the efficient approximation of functions $f(\mathbf{x}, \mathbf{y}) \in L^2(\Omega_1 \times \Omega_2)$ by either the truncated singular value decomposition

$$f_M(\mathbf{x}, \mathbf{y}) := \sum_{\ell=1}^M \sqrt{\lambda_\ell} \varphi_\ell(\mathbf{x}) \psi_\ell(\mathbf{y})$$

or by the generalized sparse grid approach

$$\widehat{f}_J(\mathbf{x}, \mathbf{y}) := \sum_{j_1/\sigma + j_2 \leq J} \sum_{k_1 \in \mathbb{V}_{j_1}^{(1)}} \sum_{k_2 \in \mathbb{V}_{j_2}^{(2)}} \beta_{(j_1, k_1), (j_2, k_2)} \xi_{j_1, k_1}^{(1)}(\mathbf{x}) \xi_{j_2, k_2}^{(2)}(\mathbf{y})$$

where $\sigma > 0$ is an appropriately chosen parameter. In the first representation, $\{\varphi_\ell\}_{\ell=1}^M$ and $\{\psi_\ell\}_{\ell=1}^M$ are sets of orthonormal functions. They are a-priorily unknown, can in general not be derived analytically

and need thus to be approximated in the ansatz spaces $\{V_j^{(i)}\}$. In other words, the approximation involves in most applications both, a truncation after M terms *and* an approximate computation of the singular values and the associated left and right singular vectors. In the second representation, $\{\xi_{j,k}^{(i)}\}_{k \in \nabla_j^{(i)}, j \in \mathbb{N}}$ are in general multilevel or wavelet bases associated with the trial spaces, where the index j refers to the level of resolution and the index k refers to the locality of the basis function (the precise definition will be given in Section 4).

For our comparison we consider the smoothness of the function f to be measured in isotropic Sobolev norms. The result of this paper is then as follows: given a function $f \in H^p(\Omega_1 \times \Omega_2)$, we have to spend $\mathcal{O}(\varepsilon^{-q})$ with $q = \frac{2 \min\{r,p\} \min\{n_1,n_2\} + 2p \max\{n_1,n_2\}}{(2p - \min\{n_1,n_2\}) \min\{r,p\}}$ degrees of freedom for the computation of the approximative truncated singular value decomposition to reach a specific prescribed accuracy ε . For the general sparse grid method with associated parameter $\sigma = \sqrt{n_1/n_2}$ (a precise definition is given in Section 4) it is known that this cost-complexity rate is of the order $\mathcal{O}(\varepsilon^{-q})$ with $q = \max\{\frac{n_1+n_2}{p}, \frac{n_1}{r}, \frac{n_2}{r}\}$ (Griebel & Harbrecht (2011)). We compare the two approaches and present numerical results which demonstrate that these rates are also achieved in practice.

Note that we have only an upper bound for the truncated singular value decomposition. Therefore, our comparison of the truncated singular value decomposition and the sparse grid approach (see Section 5) needs to be interpreted carefully. Indeed, the present paper is only a first attempt towards a “fair” comparison of both approaches, which especially takes into account that the eigenvectors of the approximative truncated singular value decomposition need to be computed and stored in any practical numerical approach.

The remainder of this article is organized as follows: In Section 2 we give a short introduction to multilevel approximation. In Section 3 we describe the singular value decomposition of a two-valued function on $\Omega_1 \times \Omega_2$ and discuss its approximation properties in detail. Section 4 gives the basics of the so-called general sparse grid approximation of a two-valued function on $\Omega_1 \times \Omega_2$ and presents its error rates and cost complexities. In Section 5, we compare the two approximations. In Section 6, we present numerical results which show that the theoretical complexity rates are also achieved in practice. Section 7 concludes with some final remarks.

Throughout this paper, the notion “essential” in connection with the complexity estimates means “up to logarithmic terms”. Moreover, to avoid the repeated use of generic but unspecified constants, we denote by $C \lesssim D$ that C is bounded by a multiple of D independently of parameters which C and D may depend on. Obviously, $C \gtrsim D$ is defined as $D \lesssim C$, and $C \sim D$ as $C \lesssim D$ and $C \gtrsim D$.

2. Approximation on the subdomains

Let $\Omega \subset \mathbb{R}^n$ be a sufficiently smooth, bounded domain. In general, one uses finite elements to approximate functions on $L^2(\Omega)$. In the present paper we focus on the common h -method, i.e., on finite elements of fixed approximation order. Then, particularly for applying multiscale techniques, one has a sequence of nested trial spaces

$$V_0 \subset V_1 \subset V_2 \subset \dots \subset L^2(\Omega) \quad (2.1)$$

such that

$$L^2(\Omega) = \overline{\bigcup_{j \in \mathbb{N}_0} V_j},$$

which is called *multiscale analysis*. Each space V_j is defined by a single scale basis $\Phi_j = \{\phi_{j,k}\}$, i.e. $V_j = \text{span}\{\phi_{j,k} : k \in \Delta_j\}$, where Δ_j denotes a suitable index set with cardinality $\#\Delta_j \sim 2^{nj}$.

We say that the trial spaces have (*approximation*) *order* $r \in \mathbb{N}$ if

$$r = \sup \left\{ s \in \mathbb{R} : \inf_{v_j \in V_j} \|v - v_j\|_{L^2(\Omega)} \lesssim h_j^s \|v\|_s \quad \forall v \in H^s(\Omega) \right\}, \quad (2.2)$$

where the quantity $h_j \sim 2^{-j}$ corresponds to the mesh width associated with the subspace V_j on Ω . Note that the integer $r > 0$ refers in general to the maximal order of polynomials which are locally contained in V_j .

Equation (2.2) implies that a given function $v \in H^p(\Omega)$, $0 \leq p \leq r$, can be approximated in V_j at a rate h_j^p , that is

$$\inf_{v_j \in V_j} \|v - v_j\|_{L^2(\Omega)} \lesssim h_j^p \|v\|_{H^p(\Omega)}, \quad 0 \leq p \leq r. \quad (2.3)$$

Thus, when we approximate a function $v \in H^p(\Omega)$ with $0 \leq p \leq r$ by uniform mesh refinement we obtain the rate h_j^p according to (2.3). Since the meshsize and the number of unknowns in V_j are related by $\dim(V_j) \sim 2^{jn} \sim h_j^{-n}$, we deduce that

$$N \sim \varepsilon^{-n/p} \quad (2.4)$$

unknowns have to be spent to achieve an approximation error ε . The best possible rate $N^{-n/r}$ is achieved if $p = r$, that is if $v \in H^r(\Omega)$.

For our subsequent analysis, to efficiently approximate two-variate functions in $L^2(\Omega_1 \times \Omega_2)$, we shall fix the definitions, properties and cost complexities individually for each subdomain $\Omega_i \in \mathbb{R}^{n_i}$, $i = 1, 2$. That is, we fix two multiscale analyses

$$V_0^{(i)} \subset V_1^{(i)} \subset V_2^{(i)} \subset \dots \subset L^2(\Omega_i), \quad i = 1, 2 \quad (2.5)$$

which are assumed to provide the *same* approximation order r . The construction of the approximative truncated singular value decomposition in Section 3 and the sparse grid in Section 4 are based on these multiscale analyses. A comparison for the case of the *same* r , i.e., where the underlying finite element spaces in (2.5) for both, the approximative truncated singular value decomposition and the sparse grid, have the same order r , then takes place in Section 5. The question we address in this paper is as follows: Given a function $H^p(\Omega_1 \times \Omega_2)$, where $p > 0$ is arbitrary and where trial spaces with fixed order r are used in both approaches, which algorithm provides the cheaper approximation?

3. Singular value decomposition

3.1 Definition and mapping properties

We intend to numerically represent functions $f(\mathbf{x}, \mathbf{y}) \in L^2(\Omega_1 \times \Omega_2)$ on tensor product domains $\Omega_1 \times \Omega_2$ in an efficient way. One way to solve this approximation problem is to use an ansatz by means of tensor products which separates the variables \mathbf{x} and \mathbf{y} . We first consider the approximation

$$f(\mathbf{x}, \mathbf{y}) \approx f_M(\mathbf{x}, \mathbf{y}) = \sum_{\ell=0}^M \alpha_\ell \varphi_\ell(\mathbf{x}) \psi_\ell(\mathbf{y}), \quad (3.1)$$

with certain coefficients $\alpha_\ell \in \mathbb{R}$ and normalized functions $\varphi_\ell \in L^2(\Omega_1)$ and $\psi_\ell \in L^2(\Omega_2)$. Such an approximation is called *low rank approximation*.

It is well known (see e.g. Ghanem & Spanos (1991), L oeve (1978), or Schwab & Todor (2006)) that, with respect to the number M of terms, the best possible representation of a function $f \in L^2(\Omega_1 \times \Omega_2)$

in the L^2 -sense is given by the Karhunen-Lòeve / singular value decomposition. Then, $\alpha_\ell = \sqrt{\lambda_\ell}$ are the eigenvalues of the below defined integral operator (3.2) with kernel (3.3). The decay of the eigenvalues is important for the fast convergence (in terms of M) of the series (3.1). As we will see in Subsection 3.2, the decay depends on the smoothness of the function f to be approximated. We analyze the truncation error in Subsection 3.3 and discuss the smoothness of the eigenfunctions in Subsection 3.4. In Subsection 3.5 we finally consider the numerical treatment of (3.1). Besides determining the coefficients $\{\alpha_\ell\}_{\ell \in \mathbb{N}}$, a numerical scheme needs to approximate the functions $\{\varphi_\ell\}_{\ell \in \mathbb{N}}$ and $\{\psi_\ell\}_{\ell \in \mathbb{N}}$ in appropriate trial spaces $V_{j_1}^{(1)}$ and $V_{j_2}^{(2)}$, respectively, up to an accuracy corresponding to that of (3.1). Recall that the trial spaces which we consider are elements of the multiscale analyses (2.5) which have the same approximation order r .

To derive the singular value decomposition, we shall consider the integral operator

$$\mathcal{S} : L^2(\Omega_1) \rightarrow L^2(\Omega_2), \quad (\mathcal{S}u)(\mathbf{y}) := \int_{\Omega_1} f(\mathbf{x}, \mathbf{y})u(\mathbf{x})d\mathbf{x}.$$

Its adjoint is

$$\mathcal{S}^* : L^2(\Omega_2) \rightarrow L^2(\Omega_1), \quad (\mathcal{S}^*u)(\mathbf{x}) := \int_{\Omega_2} f(\mathbf{x}, \mathbf{y})u(\mathbf{y})d\mathbf{y}.$$

Then, to obtain the low-rank representation (3.1) we need to compute the eigenvalues of the integral operator

$$\mathcal{K} = \mathcal{S}^*\mathcal{S} : L^2(\Omega_1) \rightarrow L^2(\Omega_1), \quad (\mathcal{K}u)(\mathbf{x}) := \int_{\Omega_1} k(\mathbf{x}, \mathbf{x}')u(\mathbf{x}')d\mathbf{x}' \quad (3.2)$$

whose kernel function is given by

$$k(\mathbf{x}, \mathbf{x}') = \int_{\Omega_2} f(\mathbf{x}, \mathbf{y})f(\mathbf{x}', \mathbf{y})d\mathbf{y} \in L^2(\Omega_1 \times \Omega_1). \quad (3.3)$$

This is a Hilbert-Schmidt kernel. Thus, the associated integral operator \mathcal{K} is compact. Moreover, since \mathcal{K} is self-adjoint, there exists a decomposition into eigenpairs $(\lambda_\ell, \varphi_\ell)$ which fulfills

$$\mathcal{K}\varphi_\ell = \lambda_\ell\varphi_\ell, \quad \ell \in \mathbb{N},$$

with non-negative eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_m \rightarrow 0$ and eigenfunctions $\{\varphi_\ell\}_{\ell \in \mathbb{N}}$, which constitute an orthonormal basis in $L^2(\Omega_1)$.

We now define for all $\ell \in \mathbb{N}$ with $\lambda_\ell > 0$ the function $\psi_\ell \in L^2(\Omega_2)$ by

$$\psi_\ell(\mathbf{y}) = \frac{1}{\sqrt{\lambda_\ell}}(\mathcal{S}\varphi_\ell)(\mathbf{y}) = \frac{1}{\sqrt{\lambda_\ell}} \int_{\Omega_1} f(\mathbf{x}, \mathbf{y})\varphi_\ell(\mathbf{x})d\mathbf{x}. \quad (3.4)$$

This constitutes a second sequence of orthonormal functions since

$$\begin{aligned} (\psi_k, \psi_\ell)_{L^2(\Omega_2)} &= \frac{1}{\sqrt{\lambda_k\lambda_\ell}}(\mathcal{S}\varphi_k, \mathcal{S}\varphi_\ell)_{L^2(\Omega_2)} = \frac{1}{\sqrt{\lambda_k\lambda_\ell}}(\mathcal{K}\varphi_k, \varphi_\ell)_{L^2(\Omega_1)} \\ &= \frac{\lambda_k}{\sqrt{\lambda_k\lambda_\ell}}(\varphi_k, \varphi_\ell)_{L^2(\Omega_1)} = \delta_{k,\ell}. \end{aligned}$$

If $\lambda_\ell = 0$ for some $\ell \in \mathbb{N}$ we can extend this collection of functions properly to obtain an orthonormal basis $\{\psi_\ell\}_{\ell \in \mathbb{N}}$ of $L^2(\Omega_2)$. Due to

$$\sqrt{\lambda_\ell} \varphi_\ell(\mathbf{x}) = \frac{1}{\sqrt{\lambda_\ell}} (\mathcal{S}^* \mathcal{S} \varphi_\ell)(\mathbf{x}) = (\mathcal{S}^* \psi_\ell)(\mathbf{x}) = \int_{\Omega_2} f(\mathbf{x}, \mathbf{z}) \psi_\ell(\mathbf{z}) d\mathbf{z} \quad (3.5)$$

for all $\mathbf{x} \in \Omega_1$ and $\ell \in \mathbb{N}$, we finally obtain the representation

$$f(\mathbf{x}, \mathbf{y}) = \sum_{\ell=0}^{\infty} \sqrt{\lambda_\ell} \varphi_\ell(\mathbf{x}) \psi_\ell(\mathbf{y}). \quad (3.6)$$

With (3.4) and (3.5), this equation is easily verified by testing with the orthonormal basis $\{\varphi_k \otimes \psi_\ell\}_{k, \ell \in \mathbb{N}}$ of $L^2(\Omega_1 \times \Omega_2)$.

REMARK 3.1 The adjoint kernel $\tilde{k}(\cdot, \cdot)$ is just obtained by interchanging Ω_1 and Ω_2 , i.e.,

$$\tilde{k}(\mathbf{y}, \mathbf{y}') = \int_{\Omega_1} f(\mathbf{x}, \mathbf{y}) f(\mathbf{x}, \mathbf{y}') d\mathbf{x} \in L^2(\Omega_2 \times \Omega_2).$$

Then, one has the integral operator

$$\tilde{\mathcal{K}} = \mathcal{S} \mathcal{S}^* : L^2(\Omega_2) \rightarrow L^2(\Omega_2), \quad (\tilde{\mathcal{K}} u)(\mathbf{y}) := \int_{\Omega_2} \tilde{k}(\mathbf{y}, \mathbf{y}') u(\mathbf{y}') d\mathbf{y}'.$$

Again there exists a decomposition into eigenpairs

$$\tilde{\mathcal{K}} \tilde{\varphi}_\ell = \tilde{\lambda}_\ell \tilde{\varphi}_\ell, \quad \ell \in \mathbb{N},$$

with non-negative eigenvalues $\tilde{\lambda}_1 \geq \tilde{\lambda}_2 \geq \dots \geq \tilde{\lambda}_m \rightarrow 0$ and eigenfunctions $\tilde{\varphi}_\ell \in L^2(\Omega_2)$. We also obtain a second sequence of orthonormal functions $\tilde{\psi}_\ell \in L^2(\Omega_1)$ analogously to (3.4). The functions $\{\tilde{\varphi}_\ell\}_{\ell \in \mathbb{N}}$ and $\{\tilde{\psi}_\ell\}_{\ell \in \mathbb{N}}$ will be the same as before but now their roles are exchanged. Moreover, the eigenvalues λ_ℓ and $\tilde{\lambda}_\ell$ of \mathcal{K} and $\tilde{\mathcal{K}}$ coincide.

We shall prove the following auxiliary result concerning the mapping properties of the integral operators \mathcal{S} and \mathcal{S}^* . To this end, for $s > 0$ we mean by $H^{-s}(\Omega) := (H^s(\Omega))'$ the dual of the Sobolev space $H^s(\Omega)$ (which is usually denoted by $\tilde{H}^{-s}(\Omega)$). Moreover, for $s_1, s_2 \in \mathbb{R}$, let

$$H_{mix}^{s_1, s_2}(\Omega_1 \times \Omega_2) := H^{s_1}(\Omega_1) \otimes H^{s_2}(\Omega_2)$$

denote the Sobolev space of dominating mixed derivatives.

LEMMA 3.1 Assume that $f \in H^p(\Omega_1 \times \Omega_2)$. Then, the operators

$$\mathcal{S} : H^{-s}(\Omega_1) \rightarrow H^{p-s}(\Omega_2), \quad \mathcal{S}^* : H^{-s}(\Omega_2) \rightarrow H^{p-s}(\Omega_1)$$

are continuous for all $s \in [0, p]$.

Proof. From $H^p(\Omega_1 \times \Omega_2) \subset H_{mix}^{0,p}(\Omega_1 \times \Omega_2)$ it follows that $f \in H_{mix}^{0,p}(\Omega_1 \times \Omega_2)$. Therefore, the operator

$\mathcal{S} : L^2(\Omega_1) \rightarrow H^p(\Omega_2)$ is continuous since

$$\begin{aligned}
\|\mathcal{S}u\|_{H^p(\Omega_2)} &= \sup_{\|v\|_{H^{-p}(\Omega_2)}=1} (\mathcal{S}u, v)_{L^2(\Omega_2)} \\
&= \sup_{\|v\|_{H^{-p}(\Omega_2)}=1} (f, u \otimes v)_{L^2(\Omega_1 \times \Omega_2)} \\
&\leq \sup_{\|v\|_{H^{-p}(\Omega_2)}=1} \|f\|_{H_{mix}^{0,p}(\Omega_1 \times \Omega_2)} \|u \otimes v\|_{H_{mix}^{0,-p}(\Omega_1 \times \Omega_2)} \\
&\sim \|f\|_{H_{mix}^{0,p}(\Omega_1 \times \Omega_2)} \|u\|_{L^2(\Omega_1)}.
\end{aligned}$$

Note that we have used here that $H_{mix}^{0,-p}(\Omega_1 \times \Omega_2) = L^2(\Omega_1) \otimes H^{-p}(\Omega_2)$.

In complete analogy one shows that $\mathcal{S}^* : L^2(\Omega_2) \rightarrow H^p(\Omega_1)$ is continuous which proves the desired assertion for $s = 0$. By duality one also infers the assertion for $s = p$. The assertion for $s \in (0, p)$ is finally obtained by interpolation. \square

3.2 Decay of the eigenvalues

With the above lemma we are now able to determine the decay rate of the eigenvalues of the integral operator $\mathcal{K} = \mathcal{S}^* \mathcal{S}$ with kernel (3.3).

THEOREM 3.1 Consider $f \in H^p(\Omega_1 \times \Omega_2)$ with associated kernel k from (3.3) and associated integral operator \mathcal{K} from (3.2). Then, the eigenvalues $\{\lambda_\ell\}_{\ell \in \mathbb{N}}$ of \mathcal{K} decay like

$$\lambda_\ell \lesssim \ell^{-\frac{2p}{\min\{n_1, n_2\}}} \quad \text{as } \ell \rightarrow \infty. \quad (3.7)$$

Proof. We shall investigate the dependence of the decay of the eigenvalues of the integral operator $\mathcal{K} = \mathcal{S}^* \mathcal{S}$ on the smoothness p of f . To this end, we introduce *new*¹ finite element spaces $U_N \subset L^2(\Omega_2)$, which consist of N discontinuous, piecewise polynomial functions of total degree $[p]$ on a quasi-uniform triangulation of Ω_2 with mesh width $h_N \sim N^{-1/n_2}$. Then, the Bramble-Hilbert lemma implies the following approximation result for the L^2 -orthogonal projection $P_N : L^2(\Omega_2) \rightarrow U_N$:

$$\|(I - P_N)w\|_{L^2(\Omega_2)} \leq c_p N^{-p/n_2} \|w\|_{H^p(\Omega_2)},$$

provided that $w \in H^p(\Omega_2)$. Then, since $\mathcal{S}^* P_N \mathcal{S} : L^2(\Omega_1) \rightarrow L^2(\Omega_1)$ is an operator of finite rank N , the

¹The present proof relies on an approximation argument. The new finite element spaces $\{U_N\}$ are introduced to obtain the optimal convergence rate with N degrees of freedom.

min-max principle of Courant-Fischer implies

$$\begin{aligned}
\lambda_{N+1} &= \min_{\substack{V \subset L^2(\Omega_1) \\ \dim V \leq N}} \max_{\substack{u \in V \\ \|u\|_{L^2(\Omega_1)}=1}} (\mathcal{H}u, u)_{L^2(\Omega_1)} \\
&\leq \max_{\substack{u \perp \text{img}(\mathcal{S}^* P_N \mathcal{S}) \\ \|u\|_{L^2(\Omega_1)}=1}} (\mathcal{S}^* \mathcal{S} u, u)_{L^2(\Omega_1)} \\
&= \max_{\substack{u \perp \text{img}(\mathcal{S}^* P_N \mathcal{S}) \\ \|u\|_{L^2(\Omega_1)}=1}} (\mathcal{S}^*(I - P_N)\mathcal{S}u, u)_{L^2(\Omega_1)} \\
&= \max_{\substack{u \perp \text{img}(\mathcal{S}^* P_N \mathcal{S}) \\ \|u\|_{L^2(\Omega_1)}=1}} ((I - P_N)\mathcal{S}u, (I - P_N)\mathcal{S}u)_{L^2(\Omega_2)} \\
&\leq \sup_{\|u\|_{L^2(\Omega_1)}=1} \|(I - P_N)\mathcal{S}u\|_{L^2(\Omega_2)}^2 \\
&\lesssim N^{-2p/n_2} \sup_{\|u\|_{L^2(\Omega_1)}=1} \|\mathcal{S}u\|_{H^p(\Omega_2)}^2.
\end{aligned}$$

Since $\mathcal{S} : L^2(\Omega_1) \rightarrow H^p(\Omega_2)$ is continuous according to Lemma 3.1, we arrive at $\lambda_{N+1} \lesssim N^{-2p/n_2}$. Applying the same arguments to the operator \mathcal{H} , one gets the decay rate $\lambda_{N+1} \lesssim N^{-2p/n_1}$. Thus, $\lambda_{N+1} \lesssim N^{-2p/\min\{n_1, n_2\}}$. Substituting finally $N + 1$ by ℓ yields the desired result. \square

REMARK 3.2 The min-max principle of Courant-Fischer is also used in Schwab & Todor (2006) to estimate the decay of the eigenvalues. There, however, only the regularity of the kernel function $k \in H_{\text{mix}}^{p,p}(\Omega_1 \times \Omega_1)$ in the first variable enters the estimate which gives the rate

$$\lambda_\ell \lesssim \ell^{-\frac{p}{\min\{n_1, n_2\}}} \quad \text{as } \ell \rightarrow \infty.$$

By using the representation $\mathcal{H} = \mathcal{S}^* \mathcal{S}$, we are able to exploit the kernel's regularity in both variables which doubles the decay rate.

3.3 Truncation error

Altogether, if $f \in H^p(\Omega_1 \times \Omega_2)$, then Theorem 3.1 implies that the coefficients $\{\sqrt{\lambda_\ell}\}$ in the expansion (3.6) of f decay like $\sqrt{\lambda_\ell} \lesssim \ell^{-p/\min\{n_1, n_2\}}$. This leads to the following theorem.

THEOREM 3.2 Let $f \in H^p(\Omega_1 \times \Omega_2)$ and $p > \min\{n_1, n_2\}/2$. Then it holds

$$\left\| f - \sum_{\ell=0}^M \sqrt{\lambda_\ell} (\phi_\ell \otimes \psi_\ell) \right\|_{L^2(\Omega_1 \times \Omega_2)} \lesssim M^{\frac{1}{2} - \frac{p}{\min\{n_1, n_2\}}}. \quad (3.8)$$

Proof. Assume without loss of generality that $n_1 \leq n_2$. Then, due to $\lambda_\ell \sim \ell^{-2p/n_1}$ from Theorem 3.1, we obtain by the orthonormality of the function sets $\{\phi_\ell\}$ and $\{\psi_\ell\}$ that

$$\left\| f - \sum_{\ell=0}^M \sqrt{\lambda_\ell} (\phi_\ell \otimes \psi_\ell) \right\|_{L^2(\Omega_1 \times \Omega_2)}^2 = \left\| \sum_{\ell=M+1}^{\infty} \sqrt{\lambda_\ell} (\phi_\ell \otimes \psi_\ell) \right\|_{L^2(\Omega_1 \times \Omega_2)}^2 = \sum_{\ell=M+1}^{\infty} \lambda_\ell \lesssim \sum_{\ell=M+1}^{\infty} \ell^{-\frac{2p}{n_1}}.$$

Since $2p/n_1 > 1$ we can estimate the sum by an integral as

$$\sum_{\ell=M+1}^{\infty} \ell^{-\frac{2p}{n_1}} \leq \int_M^{\infty} x^{-\frac{2p}{n_1}} dx = \frac{M^{1-\frac{2p}{n_1}}}{\frac{2p}{n_1} - 1}$$

which leads to the desired result (3.8). \square

Consequently, to ensure the error bound

$$\left\| f - \sum_{\ell=0}^M \sqrt{\lambda_\ell} (\varphi_\ell \otimes \psi_\ell) \right\|_{L^2(\Omega_1 \times \Omega_2)} \lesssim \varepsilon \quad (3.9)$$

we need to choose the expansion degree M as

$$M \sim \varepsilon^{\frac{2 \min\{n_1, n_2\}}{\min\{n_1, n_2\} - 2p}}. \quad (3.10)$$

REMARK 3.3 The following comments are in order.

1. Regularity in terms of mixed derivatives does not further improve the results. The property $f \in H_{mix}^{p,p}(\Omega_1 \times \Omega_2)$ again yields the estimate $\lambda_\ell \lesssim \ell^{-2p/\max\{n_1, n_2\}}$ for the eigenvalues of \mathcal{K} .
2. The use of the Sobolev regularity might give a too low decay rate (3.7). For example, for the exponential kernel $k(x, y) = \exp(-|x - y|)$ on the unit square, we have $k \in H^{3/2-\delta}((0, 1) \times (0, 1))$ for all $\delta > 0$ but we observe $\sqrt{\lambda_\ell} \sim \ell^{-2}$ instead of $\sqrt{\lambda_\ell} \sim \ell^{-3/2}$, see Figure 5 in Section 6. Here, we expect that the use of Besov regularity in the approximation argument in the proof of Theorem (3.1) would give the correct decay since it allows for an adaptive refinement at the diagonal $x = y$.
3. Our considerations and thus estimate (3.8) do not apply if $0 \leq p \leq \min\{n_1, n_2\}/2$. However, it still holds $\sum_{\ell=0}^{\infty} \lambda_\ell < \infty$ since \mathcal{K} is a Hilbert-Schmidt operator.

3.4 Smoothness of the eigenfunctions

Depending on the smoothness of f , we are able to prove the following result on the regularity of the functions in the collections $\{\varphi_\ell\}$ and $\{\psi_\ell\}$. This result will be essential for any numerical computation of the truncated singular value decomposition.

LEMMA 3.2 Let $f \in H^p(\Omega_1 \times \Omega_2)$. Then, the eigenfunctions $\{\varphi_\ell\}$ and $\{\psi_\ell\}$ are in $H^p(\Omega_1)$ and $H^p(\Omega_2)$, respectively, and satisfy

$$\|\varphi_\ell\|_{H^p(\Omega_1)} \lesssim \frac{1}{\sqrt{\lambda_\ell}}, \quad \|\psi_\ell\|_{H^p(\Omega_2)} \lesssim \frac{1}{\sqrt{\lambda_\ell}}, \quad \ell \in \mathbb{N}. \quad (3.11)$$

Proof. In view of (3.5) and Lemma 3.1, we deduce

$$\|\varphi_\ell\|_{H^p(\Omega_1)} = \frac{1}{\sqrt{\lambda_\ell}} \|\mathcal{S}^* \psi_\ell\|_{H^p(\Omega_1)} \lesssim \frac{1}{\sqrt{\lambda_\ell}} \|\psi_\ell\|_{L^2(\Omega_2)} = \frac{1}{\sqrt{\lambda_\ell}}$$

for all $\ell \in \mathbb{N}$. The second estimate is shown in complete analogy. \square

3.5 Numerical approximation

So far, we used an *exact* description of the eigenfunctions. However this does not hold in practice. Instead, the eigenvalues $\{\lambda_\ell\}_{\ell=1}^M$ and eigenfunctions $\{\varphi_\ell\}_{\ell=1}^M$ and $\{\psi_\ell\}_{\ell=1}^M$ need to be approximately computed in the finite element spaces which have been introduced in Section 2.

Recall that, on both subdomains, the finite element spaces $V_j^{(i)} \subset L^2(\Omega_i)$ provide the same approximation order r , which results in the same approximation property (2.3), but, depending on the subdomains' spatial dimensions n_1, n_2 , in different cost complexities (2.4).

According to (2.3) and (3.11), when we spend $N_\varphi \sim 2^{jn_1}$ degrees of freedom for an approximation of φ_ℓ in $V_j^{(1)}$ and $N_\psi \sim 2^{jn_2}$ degrees of freedom for an approximation of ψ_ℓ in $V_j^{(2)}$, we conclude

$$\begin{aligned} \|\varphi_\ell - \varphi_{\ell,N}\|_{L^2(\Omega_1)} &\lesssim N_\varphi^{-\frac{\min\{p,r\}}{n_1}} \|\varphi_\ell\|_{H^{\min\{p,r\}}(\Omega)} \lesssim \frac{1}{\sqrt{\lambda_\ell}} N_\varphi^{-\frac{\min\{p,r\}}{n_1}}, \\ \|\psi_\ell - \psi_{\ell,N}\|_{L^2(\Omega_2)} &\lesssim N_\psi^{-\frac{\min\{p,r\}}{n_2}} \|\psi_\ell\|_{H^{\min\{p,r\}}(\Omega)} \lesssim \frac{1}{\sqrt{\lambda_\ell}} N_\psi^{-\frac{\min\{p,r\}}{n_2}}. \end{aligned} \quad (3.12)$$

Here, $\varphi_{\ell,N}$ and $\psi_{\ell,N}$ denote the numerical approximations to φ_ℓ and ψ_ℓ , respectively.

According to (3.12), to ensure $\|\varphi_\ell - \varphi_{\ell,N}\|_{L^2(\Omega_1)} \lesssim \varepsilon / \sqrt{\lambda_\ell M}$ (we will later see that this is the proper accuracy), we have to spend (cf. (3.10))

$$N_\varphi \sim \left(\frac{\varepsilon}{\sqrt{M}} \right)^{-\frac{n_1}{\min\{r,p\}}} \sim \varepsilon^{\frac{2pn_1}{(\min\{n_1,n_2\}-2p)\min\{r,p\}}}$$

unknowns for the representation of $\varphi_{\ell,N}$ and, to ensure $\|\psi_\ell - \psi_{\ell,N}\|_{L^2(\Omega_2)} \lesssim \varepsilon / \sqrt{\lambda_\ell M}$, we have to spend

$$N_\psi \sim \left(\frac{\varepsilon}{\sqrt{M}} \right)^{-\frac{n_2}{\min\{r,p\}}} \sim \varepsilon^{\frac{2pn_2}{(\min\{n_1,n_2\}-2p)\min\{r,p\}}}$$

unknowns for the representation of $\psi_{\ell,N}$, respectively. In the sequel, we will spend always $N := \max\{N_\varphi, N_\psi\}$ degrees of freedom which does not deteriorate the cost complexity. In particular, N does not depend on ℓ , i.e., all eigenfunctions $\{\varphi_{N,\ell}\}$ and $\{\psi_{N,\ell}\}$ are approximated in the same ansatz spaces.

REMARK 3.4 If $p > r$, then we even may estimate

$$\|\varphi_\ell\|_{H^r(\Omega)} \lesssim \left(\frac{1}{\sqrt{\lambda_\ell}} \right)^{\frac{r}{p}}, \quad \|\psi_\ell\|_{H^r(\Omega)} \lesssim \left(\frac{1}{\sqrt{\lambda_\ell}} \right)^{\frac{r}{p}}$$

by using interpolation arguments. Hence, (3.12) can be improved by

$$\begin{aligned} \|\varphi_\ell - \varphi_{\ell,N}\|_{L^2(\Omega_1)} &\lesssim \lambda_\ell^{-\frac{\min\{r,p\}}{2p}} N^{-\frac{\min\{p,r\}}{n_1}}, \\ \|\psi_\ell - \psi_{\ell,N}\|_{L^2(\Omega_2)} &\lesssim \lambda_\ell^{-\frac{\min\{r,p\}}{2p}} N^{-\frac{\min\{p,r\}}{n_2}}. \end{aligned}$$

As a consequence, if $p > r$, the number of unknowns for approximating the eigenfunctions can be reduced when ℓ increases. However, to exploit this fact for the computation of the truncated singular value decomposition, one needs the specific information on the smoothness index p .

We assume that the approximate eigenfunctions are normalized and pairwise orthogonal, i.e., for $1 \leq \ell, \ell' \leq M$ we have

$$\int_{\Omega_1} \boldsymbol{\varphi}_{\ell,N}(\mathbf{x}) \boldsymbol{\varphi}_{\ell',N}(\mathbf{x}) d\mathbf{x} = \delta_{\ell,\ell'}, \quad \int_{\Omega_2} \boldsymbol{\psi}_{\ell,N}(\mathbf{y}) \boldsymbol{\psi}_{\ell',N}(\mathbf{y}) d\mathbf{y} = \delta_{\ell,\ell'}.$$

Concerning the approximation of the M largest eigenvalues in the space V_N , defined via their Rayleigh quotients

$$\lambda_{\ell,N} = \int_{\Omega_1} \int_{\Omega_1} k(\mathbf{x}, \mathbf{x}') \boldsymbol{\varphi}_{\ell,N}(\mathbf{x}) \boldsymbol{\varphi}_{\ell,N}(\mathbf{x}') d\mathbf{x} d\mathbf{x}', \quad \ell = 1, 2, \dots, M.$$

we assume the following estimate

$$0 \leq \lambda_\ell - \lambda_{\ell,N} \lesssim \lambda_\ell \|\boldsymbol{\varphi}_\ell - \boldsymbol{\varphi}_{\ell,N}\|_{L^2(\Omega_1)}^2, \quad \ell = 1, 2, \dots, M. \quad (3.13)$$

We emphasize that these assumptions, in particular (3.13), are all satisfied if one computes the approximation $\{(\lambda_{N,\ell}, \boldsymbol{\varphi}_{N,\ell})\}_{\ell=1}^M$ to the eigenpairs $\{(\lambda_\ell, \boldsymbol{\varphi}_\ell)\}_{\ell=1}^M$ with a Ritz-Galerkin method in the space $V_j^{(1)}$ with $\dim V_j^{(1)} \sim N$, i.e., with $j \sim \log N$. For the details we refer the reader to Dyakonov (1996).

Based on (3.8), (3.12) and (3.13), we can now estimate the error of the discretized truncated singular value decomposition.

THEOREM 3.3 Let $f \in H^p(\Omega_1 \times \Omega_2)$ and choose

$$M \sim \varepsilon^{\frac{2 \min\{n_1, n_2\}}{\min\{n_1, n_2\} - 2p}}, \quad N \sim \varepsilon^{\frac{2p \max\{n_1, n_2\}}{(\min\{n_1, n_2\} - 2p) \min\{r, p\}}}.$$

Then, the truncated singular value decomposition satisfies the error estimate

$$\left\| f - \sum_{\ell=0}^M \sqrt{\lambda_{\ell,N}} (\boldsymbol{\varphi}_{\ell,N} \otimes \boldsymbol{\psi}_{\ell,N}) \right\|_{L^2(\Omega_1 \times \Omega_2)} \lesssim \varepsilon$$

uniformly in $\varepsilon > 0$.

Proof. It holds

$$\begin{aligned} E &:= \left\| \sum_{\ell=0}^{\infty} \sqrt{\lambda_\ell} (\boldsymbol{\varphi}_\ell \otimes \boldsymbol{\psi}_\ell) - \sum_{\ell=0}^M \sqrt{\lambda_{\ell,N}} (\boldsymbol{\varphi}_{\ell,N} \otimes \boldsymbol{\psi}_{\ell,N}) \right\|_{L^2(\Omega_1 \times \Omega_2)} \\ &\leq \left\| \sum_{\ell=M+1}^{\infty} \sqrt{\lambda_\ell} (\boldsymbol{\varphi}_\ell \otimes \boldsymbol{\psi}_\ell) \right\|_{L^2(\Omega_1 \times \Omega_2)} + \left\| \sum_{\ell=0}^M \sqrt{\lambda_\ell} (\boldsymbol{\varphi}_\ell \otimes \boldsymbol{\psi}_\ell) - \sqrt{\lambda_{\ell,N}} (\boldsymbol{\varphi}_{\ell,N} \otimes \boldsymbol{\psi}_{\ell,N}) \right\|_{L^2(\Omega_1 \times \Omega_2)}. \end{aligned}$$

According to (3.8) and (3.9) the truncation error, i.e., the first term, is bounded by $c\varepsilon$. Moreover, we split the second term into appropriate differences:

$$\begin{aligned} \sqrt{\lambda_\ell} (\boldsymbol{\varphi}_\ell \otimes \boldsymbol{\psi}_\ell) - \sqrt{\lambda_{\ell,N}} (\boldsymbol{\varphi}_{\ell,N} \otimes \boldsymbol{\psi}_{\ell,N}) &= \left(\sqrt{\lambda_\ell} - \sqrt{\lambda_{\ell,N}} \right) (\boldsymbol{\varphi}_\ell \otimes \boldsymbol{\psi}_\ell) + \sqrt{\lambda_{\ell,N}} (\boldsymbol{\varphi}_\ell \otimes (\boldsymbol{\psi}_\ell - \boldsymbol{\psi}_{\ell,N})) \\ &\quad + \sqrt{\lambda_{\ell,N}} ((\boldsymbol{\varphi}_\ell - \boldsymbol{\varphi}_{\ell,N}) \otimes \boldsymbol{\psi}_{\ell,N}). \end{aligned}$$

Thus, we arrive at

$$E \lesssim \varepsilon + \left\| \sum_{\ell=0}^M (\sqrt{\lambda_\ell} - \sqrt{\lambda_{\ell,N}}) (\varphi_\ell \otimes \psi_\ell) \right\|_{L^2(\Omega_1 \times \Omega_2)} + \left\| \sum_{\ell=0}^M \sqrt{\lambda_{\ell,N}} (\varphi_\ell \otimes (\psi_\ell - \psi_{\ell,N})) \right\|_{L^2(\Omega_1 \times \Omega_2)} \\ + \left\| \sum_{\ell=0}^M \sqrt{\lambda_{\ell,N}} ((\varphi_\ell - \varphi_{\ell,N}) \otimes \psi_{\ell,N}) \right\|_{L^2(\Omega_1 \times \Omega_2)}.$$

We now estimate the three terms on the right hand side separately. In view of (3.13), we have $\sqrt{\lambda_\ell} \geq \sqrt{\lambda_{\ell,N}}$ and, with (3.12), we obtain

$$0 \leq \left| \sqrt{\lambda_\ell} - \sqrt{\lambda_{\ell,N}} \right|^2 \leq \lambda_\ell - \lambda_{\ell,N} \lesssim \lambda_\ell \|\varphi_\ell - \varphi_{\ell,N}\|_{L^2(\Omega_1)}^2 \lesssim \frac{\varepsilon^2}{M}$$

for all $0 \leq \ell \leq M$. This yields

$$\left\| \sum_{\ell=0}^M (\sqrt{\lambda_\ell} - \sqrt{\lambda_{\ell,N}}) (\varphi_\ell \otimes \psi_\ell) \right\|_{L^2(\Omega_1 \times \Omega_2)}^2 = \sum_{\ell=0}^M \left| \sqrt{\lambda_\ell} - \sqrt{\lambda_{\ell,N}} \right|^2 \lesssim \sum_{\ell=0}^M \frac{\varepsilon^2}{M} \lesssim \varepsilon^2. \quad (3.14)$$

Next, with $\lambda_{\ell,N} \leq \lambda_\ell$, we have

$$\left\| \sum_{\ell=0}^M \sqrt{\lambda_{\ell,N}} (\varphi_\ell \otimes (\psi_\ell - \psi_{\ell,N})) \right\|_{L^2(\Omega_1 \times \Omega_2)}^2 = \sum_{\ell=0}^M \lambda_{\ell,N} \|\varphi_\ell \otimes (\psi_\ell - \psi_{\ell,N})\|_{L^2(\Omega_1 \times \Omega_2)}^2 \\ \leq \sum_{\ell=0}^M \lambda_\ell \|\varphi_\ell\|_{L^2(\Omega_1)}^2 \|\psi_\ell - \psi_{\ell,N}\|_{L^2(\Omega_2)}^2 \\ \lesssim \sum_{\ell=0}^M \frac{\varepsilon^2}{M} \lesssim \varepsilon^2, \quad (3.15)$$

and likewise

$$\left\| \sum_{\ell=0}^M \sqrt{\lambda_{\ell,N}} ((\varphi_\ell - \varphi_{\ell,N}) \otimes \psi_{\ell,N}) \right\|_{L^2(\Omega_1 \times \Omega_2)}^2 \lesssim \varepsilon^2. \quad (3.16)$$

Plugging (3.14)–(3.16) into the above estimate of E finally leads to the desired estimate $E \lesssim \varepsilon$. \square

Altogether, since we have to deal with M eigenfunctions with N coefficients each, we arrive at the following theorem.

THEOREM 3.4 The number of degrees of freedom needed to approximate a function $f \in H^p(\Omega_1 \times \Omega_2)$ by the singular value decomposition approach (3.1) with (3.12) and (3.13) to a prescribed accuracy ε is

$$\text{dof}_{\text{svd}}(\varepsilon) = M \cdot N \sim \varepsilon^{-\frac{2 \min\{r,p\} \min\{n_1,n_2\} + 2p \max\{n_1,n_2\}}{(2p - \min\{n_1,n_2\}) \min\{r,p\}}} \quad (3.17)$$

We emphasize that the estimate (3.17) does not include the work to be spent for *computing* the singular values nor the eigenfunctions. Here, a naive approach would result in a cost complexity of order

$M \cdot N^2$, the use of fast methods for nonlocal operators would result in an almost linear or even linear complexity per eigenpair. Note that in any case at least linear complexity $\mathcal{O}(M \cdot N)$ is required, which is indeed achieved with modern algorithms, see e.g. Dahmen *et al.* (2008), Dai *et al.* (2008), and Schwab & Todor (2006). Therefore, our analysis is based on the best possible situation for the approximative truncated singular value decomposition and our later comparison will be fair in this respect.

4. Sparse grids

Based on the multiscale analyses (2.5) on each individual subdomain, one naturally obtains a second method to approximate functions in tensor product spaces: By choosing complementary spaces

$$W_j^{(i)} = \text{span} \{ \xi_{j,k}^{(i)} : k \in \nabla_j^{(i)} := \Delta_j^{(i)} \setminus \Delta_{j-1}^{(i)} \}, \quad i = 1, 2,$$

such that

$$V_j^{(i)} = W_j^{(i)} \oplus V_{j-1}^{(i)}, \quad V_0^{(i)} = W_0^{(i)},$$

we can define the so called *general sparse grid space*, see Bungartz & Griebel (2004) and Griebel & Harbrecht (2011),

$$\widehat{\mathbf{V}}_J^\sigma := \bigoplus_{j_1 \sigma + j_2 / \sigma \leq J} W_{j_1}^{(1)} \otimes W_{j_2}^{(2)} \quad (4.1)$$

where $\sigma > 0$ is a given parameter. Thus, a function $\widehat{f}_J \in \widehat{\mathbf{V}}_J^\sigma$ is represented as

$$\widehat{f}_J(\mathbf{x}, \mathbf{y}) = \sum_{j_1 \sigma + j_2 / \sigma \leq J} \sum_{k_1 \in \nabla_{j_1}^{(1)}} \sum_{k_2 \in \nabla_{j_2}^{(2)}} \beta_{(j_1, k_1), (j_2, k_2)} \xi_{j_1, k_1}^{(1)}(\mathbf{x}) \xi_{j_2, k_2}^{(2)}(\mathbf{y}). \quad (4.2)$$

Sparse grids can be constructed via hierarchical bases, interpolets and wavelet-like bases (see e.g. DeVore *et al.* (1998), Griebel & Knappek (2009), Strömberg (1998), and Zenger (1991)) or even directly by finite elements in terms of frames (see e.g. Griebel (1994), Griebel & Oswald (1994), and Harbrecht *et al.* (2008b)). For a survey on sparse grids we refer the reader to Bungartz & Griebel (2004) and the references therein.

The dimension of the general sparse grid space $\widehat{\mathbf{V}}_J^\sigma$ is essentially equal to the dimension of the finest univariate finite element spaces which enter its construction, i.e., it is essentially equal to the value of $\max \{ \dim V_{J/\sigma}^{(1)}, \dim V_{J\sigma}^{(2)} \}$. Nevertheless, by considering smoothness in terms of mixed Sobolev spaces, its approximation power is essentially the same as in the full tensor product space. Precisely, we have

THEOREM 4.1 (Griebel & Harbrecht (2011)) The sparse grid space $\widehat{\mathbf{V}}_J^\sigma$ possesses

$$\dim \widehat{\mathbf{V}}_J^\sigma \sim \begin{cases} 2^{J \max\{n_1/\sigma, n_2\sigma\}}, & \text{if } n_1/\sigma \neq n_2\sigma, \\ 2^{J n_2\sigma} J, & \text{if } n_1/\sigma = n_2\sigma \end{cases}$$

degrees of freedom. Moreover, for a given function $f \in H_{\text{mix}}^{s_1, s_2}(\Omega_1 \times \Omega_2)$ with $0 < s_1, s_2 \leq r$, there holds the approximation estimate

$$\inf_{\widehat{f}_J \in \widehat{\mathbf{V}}_J^\sigma} \|f - \widehat{f}_J\|_{L^2(\Omega_1 \times \Omega_2)} \lesssim \begin{cases} 2^{-J \min\{s_1/\sigma, s_2\sigma\}} \|f\|_{H_{\text{mix}}^{s_1, s_2}(\Omega_1 \times \Omega_2)}, & \text{if } s_1/\sigma \neq s_2\sigma, \\ 2^{-J s_1/\sigma} \sqrt{J} \|f\|_{H_{\text{mix}}^{s_1, s_2}(\Omega_1 \times \Omega_2)}, & \text{if } s_1/\sigma = s_2\sigma. \end{cases}$$

In this theorem, the convergence rate is given for a function $f \in H_{mix}^{s_1, s_2}(\Omega_1 \times \Omega_2)$. In the following, however, we are interested in the convergence rate if the smoothness of f is measured in the isotropic Sobolev space $H^p(\Omega_1 \times \Omega_2)$. Since for all $s_1 + s_2 = p$ it holds $H_{mix}^{s_1, s_2}(\Omega_1 \times \Omega_2) \subset H^p(\Omega_1 \times \Omega_2)$, we have to balance s_1 and s_2 carefully if we want to compute the convergence rate of an $f \in H^p(\Omega_1 \times \Omega_2)$ in the sparse grid space $\widehat{\mathbf{V}}_J^\sigma$. The analysis given in Griebel & Harbrecht (2011) shows that the best space for approximating functions in isotropic Sobolev spaces is obtained for the choice $\sigma = \sqrt{n_1/n_2}$. With this choice, the cost complexity to approximate a function $f \in H^p(\Omega_1 \times \Omega_2)$ is given as follows.

THEOREM 4.2 (Griebel & Harbrecht (2011)) The number of degrees of freedom needed to approximate a function $f \in H^p(\Omega_1 \times \Omega_2)$ in the sparse grid space $\widehat{\mathbf{V}}_J^\sigma$ with $\sigma = \sqrt{n_1/n_2}$ to a prescribed accuracy ε is essentially

$$\text{dof}_{sg}(\varepsilon) = N \sim \varepsilon^{-\max\{\frac{n_1+n_2}{p}, \frac{n_1}{r}, \frac{n_2}{r}\}}. \quad (4.3)$$

REMARK 4.1 The sparse grid space $\widehat{\mathbf{V}}_J^\sigma$ has substantially less unknowns than the corresponding full tensor product space

$$\mathbf{V}_J^\sigma := \bigoplus_{j_1 \sigma, j_2 / \sigma \leq J} W_{j_1}^{(1)} \otimes W_{j_2}^{(2)}.$$

In this space, a function $f \in H^p(\Omega_1 \times \Omega_2)$ is approximated by the rate

$$\inf_{f_J \in \mathbf{V}_J^\sigma} \|f - f_J\|_{L^2(\Omega_1 \times \Omega_2)} \lesssim \{2^{-J \min\{p, r\}/\sigma} + 2^{-J \min\{p, r\}\sigma}\} \|f\|_{H^{\min\{p, r\}}(\Omega_1 \times \Omega_2)}$$

at the cost $\dim \mathbf{V}_J^\sigma \sim 2^{J(n_1/\sigma + n_2\sigma)}$. The choice for $\sigma = 1$ yields thus the cost complexity $N^{-\min\{p, r\}/(n_1+n_2)}$, i.e., $N \lesssim \varepsilon^{-(n_1+n_2)/\min\{p, r\}}$, which is known to be Kolmogorov's n -width for Sobolev balls in the space $H^{\min\{p, r\}}(\Omega_1 \times \Omega_2)$, see Kolmogorov (1936). Indeed, up to logarithmic terms, the sparse grid space $\widehat{\mathbf{V}}_J^\sigma$ with $\sigma = \sqrt{n_1/n_2}$ also achieves this rate. Moreover, in the case $(n_1 + n_2)/p < \max\{n_1, n_2\}/r$, it even achieves a higher rate.

5. Comparison of the two approximations

Now, for given and fixed approximation order r and spatial dimensions n_1 and n_2 , we will compare the two different approximation schemes when approximating a function $f \in H^p(\Omega_1 \times \Omega_2)$. We are interested for which values of p , the sparse grid approach is asymptotically superior to the approximative singular value decomposition. To this end, we shall distinguish three regimes of the smoothness parameter p where one should have in mind the condition $p \geq \min\{n_1, n_2\}$. The subsequent discussion has to be carefully interpreted since we deal with upper bounds only, cf. Remark 3.3.

5.1 The case $p \geq r \frac{n_1+n_2}{\max\{n_1, n_2\}}$

In this case it holds according to Theorem 4.2 that the sparse grid approach produces essentially the highest possible approximation rate $N^{-r/\min\{n_1, n_2\}}$, whereas the approximative truncated singular value decomposition reaches this rate only if f is analytical. The latter follows from the fact that $M \sim |\log(\varepsilon)|^{\min\{n_1, n_2\}}$ dominating singular values are needed, see Schwab & Todor (2006). Thus the sparse grid approach is clearly superior.

5.2 *The case $r \frac{n_1+n_2}{\max\{n_1, n_2\}} > p \geq r$*

According to Theorems 3.4 and 4.2, the truncated singular value decomposition has the complexity

$$\text{dof}_{\text{svd}}(\varepsilon) \sim \varepsilon^{-\frac{2r \min\{n_1, n_2\} + 2p \max\{n_1, n_2\}}{(2p - \min\{n_1, n_2\})r}}$$

and the sparse grid approach has the complexity

$$\text{dof}_{\text{sg}}(\varepsilon) \sim \varepsilon^{-\frac{n_1+n_2}{p}}.$$

The sparse grid approximation is asymptotically superior to the truncated singular value decomposition if $\text{dof}_{\text{sg}}(\varepsilon) \lesssim \text{dof}_{\text{svd}}(\varepsilon)$ which holds if

$$\frac{n_1 + n_2}{p} \leq \frac{2r \min\{n_1, n_2\} + 2p \max\{n_1, n_2\}}{(2p - \min\{n_1, n_2\})r}.$$

One readily infers that this inequality is equivalent to

$$0 \leq p^2 - rp + r \frac{\min\{n_1, n_2\}(n_1 + n_2)}{2 \max\{n_1, n_2\}} =: g(p).$$

The polynomial $g(p)$ might be bounded from below by

$$g(p) \geq rp - rp + r \frac{\min\{n_1, n_2\}(n_1 + n_2)}{2 \max\{n_1, n_2\}} \geq 0$$

due to $p \geq r$. Thus, the sparse grid approach exhibits a higher rate of convergence.

5.3 *The case $r > p \geq 0$*

Theorems 3.4 and 4.2 lead to the complexity

$$\text{dof}_{\text{svd}}(\varepsilon) \sim \varepsilon^{-\frac{2(n_1+n_2)}{2p - \min\{n_1, n_2\}}}$$

for the singular value decomposition and to the complexity

$$\text{dof}_{\text{sg}}(\varepsilon) \sim \varepsilon^{-\frac{n_1+n_2}{p}}$$

for the sparse grid approach. Now, it holds $\text{dof}_{\text{sg}}(\varepsilon) \lesssim \text{dof}_{\text{svd}}(\varepsilon)$ if the inequality

$$\frac{n_1 + n_2}{p} \leq \frac{2(n_1 + n_2)}{2p - \min\{n_1, n_2\}}$$

is fulfilled. As one easily checks, this inequality is indeed always satisfied. Thus, the sparse grid approach exhibits again a higher rate of convergence.

Let us remark here that in the situation $p \leq r$ an approximation in the *full* tensor product space \mathbf{V}_f^σ with $\sigma = \sqrt{n_1/n_2}$ is of the same complexity as the approximation by the sparse grid space $\widehat{\mathbf{V}}_f^\sigma$.

Altogether, we clearly see from these three cases that the general sparse grid approach is approximately at least as good as the approximative truncated singular value decomposition and is better in most situations (i.e., for an f which is less regular).

6. Numerical results

First, we present an analytical example where the sparse grid approach is superior to the truncated singular value decomposition. Afterwards, we give the results of two numerical experiments for the exponential kernel and Gaussian kernel which are important two-point correlation kernels from stochastic applications. In these examples, we thus consider the special situation that f is a symmetric function (i.e., $f(\mathbf{x}, \mathbf{y}) = f(\mathbf{y}, \mathbf{x})$) and $\Omega := \Omega_1 = \Omega_2 \subset \mathbb{R}^n$ with $n := n_1 = n_2$). Then, the singular value decomposition is simply the spectral decomposition.

6.1 An analytic example

Let f be defined as

$$f(\mathbf{x}, \mathbf{y}) = \sum_{j=0}^{\infty} \sum_{k \in \nabla_j} a_{j,k} \psi_{j,k}(\mathbf{x}) \psi_{j,k}(\mathbf{y})$$

where $\{\psi_{j,k}\}$ denotes an L^2 -normalized orthogonal wavelet basis on the domain Ω . Due to the inverse estimate we have

$$\|\psi_{j,k}\|_{H^p(\Omega)} \sim 2^{jp} \quad \text{for all } p < \gamma \quad (6.1)$$

where γ denotes the regularity of the wavelets. Consequently, the function f is in $H^p(\Omega \times \Omega)$ with $0 \leq p < \gamma$ provided that

$$\sum_{j=0}^{\infty} \sum_{k \in \nabla_j} 2^{2jp} a_{j,k}^2 < \infty. \quad (6.2)$$

The kernel (3.3) of the integral operator \mathcal{K} is given by

$$\begin{aligned} k(\mathbf{x}, \mathbf{x}') &= \sum_{j,j'=0}^{\infty} \sum_{k \in \nabla_j} \sum_{k' \in \nabla_{j'}} a_{j,k} a_{j',k'} \psi_{j,k}(\mathbf{x}) \left(\int_{\Omega} \psi_{j,k}(\mathbf{y}) \psi_{j',k'}(\mathbf{y}) d\mathbf{y} \right) \psi_{j',k'}(\mathbf{x}') \\ &= \sum_{j=0}^{\infty} \sum_{k \in \nabla_j} a_{j,k}^2 \psi_{j,k}(\mathbf{x}) \psi_{j,k}(\mathbf{x}'). \end{aligned}$$

Hence, the associated eigenvalues and eigenfunctions of \mathcal{K} are $\{a_{j,k}^2\}$ and $\{\psi_{j,k}\}$, respectively.

Estimate (6.2) implies

$$2^{2jp} a_{j,k}^2 < c 2^{-jn}, \quad k \in \nabla_j, \quad j \in \mathbb{N}_0,$$

since otherwise we would have

$$\sum_{j=0}^{\infty} \sum_{k \in \nabla_j} 2^{2jp} a_{j,k}^2 \gtrsim \sum_{j=0}^{\infty} 2^{jn} 2^{-jn} = \infty$$

due to $\#(\nabla_j) \sim 2^{jn}$. Consequently, the eigenvalues $\{a_{j,k}^2\}$ of \mathcal{K} decay essentially like

$$a_{j,k}^2 < c 2^{-j(2p+n)}, \quad k \in \nabla_j, \quad j \in \mathbb{N}_0$$

which is faster than

$$a_{j,k}^2 < c 2^{-2jp}, \quad k \in \nabla_j, \quad j \in \mathbb{N}_0$$

as predicted by Theorem 3.1. To this end, we exploited

$$2^{-jn\frac{2p}{n}} \gtrsim \ell^{-\frac{2p}{n}} \gtrsim 2^{-(j+1)n\frac{2p}{n}}.$$

As a consequence, the eigenfunctions oscillate less strongly than shown in Lemma 3.2. In fact, we have (6.1) instead of $\|\Psi_{j,k}\|_{H^p(\Omega)} \lesssim 1/\sqrt{\lambda_{j,k}} \sim 2^{j(p+n/2)}$ (cf. (3.11)). However, by choosing $p \rightarrow \gamma$, we see that the estimates in Section 3 related with the eigenvalues and eigenfunctions are sharp except for the factor $n/2$.

According to (6.2), we deduce

$$\sum_{j=J}^{\infty} \sum_{k \in \mathbb{V}_j} a_{j,k}^2 \lesssim 2^{-2Jp} \sum_{j=J}^{\infty} \sum_{k \in \mathbb{V}_j} 2^{2Jp} a_{j,k}^2 \lesssim 2^{-2Jp} \sum_{j=0}^{\infty} \sum_{k \in \mathbb{V}_j} 2^{2jp} a_{j,k}^2 \lesssim 2^{-2Jp},$$

which implies that the truncated singular value decomposition based on the first 2^{Jn} singular values approximates f with the error $\varepsilon_J \sim 2^{-Jp}$ in the L^2 -norm.

We apply properly chosen ansatz functions of approximation power r to approximate the continuous eigenfunctions $\{\psi_{j,k}\}$ by their discrete counterparts $\{\Psi_{j,k,N}\}$. Thus, in view of (2.3) and by using (6.1) with $p = \min\{\gamma, r\}$, we have

$$\|\Psi_{j,k} - \Psi_{j,k,N}\|_{L^2(\Omega)} \lesssim N_j^{-\min\{\gamma,r\}/n} 2^{j\min\{\gamma,r\}}.$$

Note that we use here an ansatz which spends the optimal number of unknowns to approximate the eigenfunction $\{\psi_{j,k}\}$, i.e., we employ N_j degrees of freedom to approximate the function $\{\psi_{j,k}\}$ instead of a fixed number N which is identical for all eigenfunctions. This is different from Section 3 and even favours the approximative truncated singular value decomposition over the sparse grid approach, cf. Remark 3.4.

To ensure the necessary accuracy $\varepsilon_J/\sqrt{M_J \lambda_{j,k}}$, we have to balance

$$\|\Psi_{j,k} - \Psi_{j,k,N}\|_{L^2(\Omega)} \stackrel{!}{\sim} \frac{\varepsilon_J}{\sqrt{M_J \lambda_{j,k}}} \sim \frac{2^{-Jp}}{2^{Jn/2} 2^{-j(p+n/2)}},$$

compare Section 3. To this end, we essentially have to spend

$$N_j \sim \left(\frac{2^{-(J-j)p}}{2^{(J-j)n/2} 2^{j\min\{\gamma,r\}}} \right)^{-n/\min\{\gamma,r\}} \sim 2^{\frac{(J-j)(p+n/2)n}{\min\{\gamma,r\}}} 2^{jn}$$

unknowns. Therefore, to represent the first 2^{Jn} eigenfunctions, we need at least (up to a constant)

$$\sum_{j=0}^J N_j 2^{jn} = \sum_{j=0}^J 2^{\frac{(J-j)(p+n/2)n}{\min\{\gamma,r\}}} 2^{2jn} = 2^{2Jn} \sum_{j=0}^J 2^{\frac{(J-j)(p+n/2-2\min\{\gamma,r\})n}{\min\{\gamma,r\}}} \quad (6.3)$$

degrees of freedom. Altogether, for our special example, this shows that the truncated and approximated singular value decomposition involves the cost complexity (6.3) to obtain the error accuracy $\varepsilon_J \sim 2^{-Jp}$.

In contrast to that, the general sparse grid approach with $\sigma = \sqrt{n_1/n_2} = 1$ produces essentially the same error rate $\varepsilon_J \sim 2^{-Jp}$ with 2^{2Jn} unknowns. If $p + n/2 > 2\min\{\gamma, r\}$, the cost complexity (6.3) is significantly larger than 2^{2Jn} . For example, choosing $p \rightarrow \gamma$ and $r > \gamma$, we get

$$n > 2\gamma$$

which describes a situation where the sparse grid approach is asymptotically superior to the corresponding approximative truncated singular value decomposition. Otherwise, the approximative truncated singular value decomposition is superior to the corresponding sparse grid approach.

In case of the singular value decomposition, we have two contradicting effects. On the one hand, the eigenfunctions can be arbitrarily smooth as for example in the case of Fourier series. Then, the smoothness is limited by the decay rate of the singular values. On the other hand, also the eigenvalue decay can be arbitrarily fast while the eigenfunctions are not smooth. A simple example are functions of finite rank but with eigenfunctions of low regularity as in the above example. Both properties affect the cost complexity of the truncated singular value decomposition differently. Compared with this, the convergence of the general sparse grid approach rather depends on the smoothness of the function to be approximated.

6.2 Gaussian kernel

In the second example let Ω be the unit interval $(0, 1)$ (i.e. $n = 1$). On level j , we subdivide $(0, 1)$ into 2^j intervals of length 2^{-j} which leads to $2^j + 1$ ansatz functions. The approximation spaces V_j under consideration are then generated by continuous piecewise linear polynomials on an equidistant partition of the interval $(0, 1)$ (i.e., $r = 2$).

We discuss the approximation of the Gaussian kernel

$$f(x, y) = \frac{1}{\sqrt{2\pi\theta}} \exp\left(-\frac{|x-y|^2}{\theta^2}\right), \quad \theta > 0.$$

Since f is known to be analytical and thus arbitrarily smooth, particularly $f \in H_{mix}^{2,2}((0, 1) \times (0, 1))$, the general sparse grid approach with $\sigma = \sqrt{n_1/n_2} = 1$ converges with the optimal rate $h_j^2 \sqrt{j} \sim 4^{-j} \sqrt{j}$. This is also observed by our numerical results, see Figure 1. Of course, the smaller the value of θ gets the larger the constant becomes which appears in front of the complexity estimate.




FIG. 1. The L^2 -error of the sparse grid approximation of the Gaussian kernel converges essentially like N^{-2} .

According to Schwab & Todor (2006), the singular values of an analytical kernel decay exponentially (if $n_1 = n_2 = 1$). In fact, in case of the Gaussian kernel the eigenvalues decay even double exponentially, cf. Figure 2, where the decay is the faster the larger θ is. Since the eigenfunctions are also analytical, it holds

$$\|\varphi_\ell\|_{H^2(0,1)} \lesssim \lambda_\ell^{-s} \quad \text{for any } s > 0.$$

Therefore, it suffices to compute the eigenfunctions with accuracy $\mathcal{O}(h_j^2)$ (see Schwab & Todor (2006) for the details) which leads to a quadratic rate of convergence with $\mathcal{O}(N \log^{1/2}(N))$ cost. This rate is also validated by Figure 3. Here, we computed the L^2 -error of the truncated and approximated singular value decomposition for all discretization levels $j = 6, 7, \dots, 15$ and truncation lengths $M = 2, 4, \dots, 40$. We observe that an increase of the truncation length improves double exponentially the approximation quality until a certain limit is reached which is induced by the approximation error of the discrete eigenfunctions. Vice versa, the increase of the discretization level gives a convergence rate $N^{-2} \sim 4^{-j}$ until a certain limit is reached which is induced by the truncation error. We thus obtain the quadratic rate $N^{-2} \sim 4^{-j}$ by proportionally increasing \sqrt{M} and $N = 2^j + 1$ which yields the above predicted cost complexity.




FIG. 2. The eigenvalues of the Gaussian kernel decay double exponentially.

Altogether, both approaches converge in our setting for the Gaussian kernel with essentially the same rate. As it is seen in Figure 4, the particular constants which are involved in the estimates of the cost complexities seem to be comparable as well. We plotted the cost complexities of the sparse grid approach (blue lines) and the truncated and approximated singular value decomposition (green lines) for different values of ε and θ . The resulting graphs are quite similar. Moreover, it is observed that the cost complexities increase for both approaches by about the same factors if θ decreases. Note that the slightly different slopes of the graphs result from the fact that the sparse grid approach produces the rate $N^{-2}\sqrt{\log N}$ at the cost $N \log N$ whereas the truncated and approximated singular value decomposition produces the rate N^{-2} at the cost $N\sqrt{\log N}$.




FIG. 3. Truncation length M and discretization level j versus accuracy in case of the Gaussian kernel.

6.3 Exponential kernel

Again, let $\Omega = (0, 1)$ and consider piecewise linear ansatz functions. Our third example is concerned with the approximation of the exponential kernel

$$f(x, y) = \exp(-|x - y|).$$

Since f is only Lipschitz continuous at the diagonal $x = y$, it follows that $f \in H^{3/2-\delta}((0, 1) \times (0, 1))$ for any $\delta > 0$. Therefore, according to Theorem 3.4, we can essentially guarantee the rate $\text{dof}_{\text{svd}}(\varepsilon) \sim \varepsilon^{-2}$. Nonetheless, as already mentioned in Remark 3.3, the singular values decay like $\sqrt{\lambda_\ell} \sim \ell^{-2}$ which is faster than predicted. In addition, it turned out in our numerical tests that also the eigenfunctions $\{\varphi_\ell\}$ are more regular than expected, i.e., they satisfy $\|\varphi_\ell\|_{H^2(0,1)} \sim 1/\sqrt{\lambda_\ell} \sim \ell^2$. The first nine eigenfunctions are depicted in Figure 6 which clearly shows the smoothness of the eigenfunctions. By repeating our analysis with these settings (that is, $p = r = 2$ and $n_1 = n_2 = 1$) one can show that

$$\text{dof}_{\text{svd}}(\varepsilon) \sim \varepsilon^{-4/3} \quad (6.4)$$

which is much better than predicted. In particular, it is the same rate as produced by a approximation on a uniform grid on $(0, 1) \times (0, 1)$. This fact is also seen in the left plot of Figure 7 where we displayed the L^2 -error of the truncated and approximated singular value decomposition for all discretization levels $j = 6, 7, \dots, 15$ and truncation lengths $M = 2^k$, $k = 0, 1, \dots, j$. The best cost complexity is offered by choosing $N \sim M \sim \varepsilon^{-2/3}$ which results in the rate (6.4).

On the other hand, in accordance with Theorem 4.2, the sparse grid approach with $\sigma = \sqrt{n_1/n_2} = 1$ realizes (essentially) the same rate: $\text{dof}_{\text{sg}}(\varepsilon) \sim \varepsilon^{-4/3}$. Indeed, this is validated by our computations (see the left plot of Figure 8).

In the left plot of Figure 9, we compare the cost complexities of the sparse grid approach (blue line) and the truncated and approximated singular value decomposition (red line). We observe that




FIG. 4. Comparison of the cost complexities in case of the Gaussian kernel.

both approaches possess essentially the rate $\varepsilon^{-4/3}$ (dashed black line). The cost complexity of the truncated and approximated singular value decomposition is identical to the cost complexity of the full tensor product approximation. Hence, it is somewhat better than the cost complexity of the sparse grid approach which involves an additional logarithmic factor.

Note here that only a conventional general sparse grid is involved and no locally adapted sparse grid is used, see e.g. Bungartz & Griebel (2004) and the references therein. For this approach, we expect a doubling of the convergence rate, i.e., essentially the rate $N^{-1.5}$. The analysis of such a nonlinear approximation scheme is however beyond of the scope of this paper.

6.4 Matérn kernel

For our last example, we choose the Matérn kernel

$$f(x, y) = (1 + |x - y|) \exp(-|x - y|)$$

and use the same numerical set-up as before. The function f is C^2 -smooth with second derivatives which are Lipschitz-continuous at the diagonal $x = y$. Hence, it follows that $f \in H^{7/2-\delta}((0, 1) \times (0, 1))$ for any $\delta > 0$. As can be seen in the right plot of Figure 5, the singular values decay now like $\sqrt{\lambda_\ell} \sim \ell^{-4}$ which is again by the factor $1/2$ better than predicted in Theorem 3.1. Moreover, from the right plot of Figure 7, by fixing the truncation length M , it can be inferred that the eigenfunctions are approximated at the rate N^{-2} . This is slightly better than to be expected from (3.12) but compare here Remark 3.4. Altogether, for the truncated and approximated singular value decomposition, we get the cost complexity $\varepsilon^{-11/14}$ by balancing $N^{-2} \sim M^{-7/2} \sim \varepsilon$. Compared with this, Theorem 3.4 gives only the rate $\varepsilon^{-4/3}$.

In comparison, the sparse grid approach with N degrees of freedom converges essentially like $N^{-7/4}$, see the right plot of Figure 8. This is in agreement with Theorem 4.2 which states the cost complexity $\text{dof}_{\text{sg}}(\varepsilon) \sim \varepsilon^{-4/7}$. Consequently, for this example, the sparse grid approach is indeed superior to the truncated and approximated singular value decomposition. This is also confirmed by the right plot of




FIG. 5. The eigenvalues of the exponential kernel (left) and the Matérn kernel (right).




FIG. 6. The first nine eigenfunctions of the exponential kernel.




FIG. 7. Truncation length M and discretization level j versus accuracy in case of the exponential kernel (left) and the Matérn kernel (right).




FIG. 8. The sparse grid approximations of the exponential kernel (left) and the Matérn kernel (right) converge essentially like $N^{-0.75}$ and $N^{-1.75}$, respectively.




FIG. 9. The cost complexities of the exponential kernel (left) and the Matérn kernel (right).

Figure 9. The cost complexity of the sparse grid approach (blue line) admits a better rate than the cost complexity of the truncated and approximated singular value decomposition (red line).

7. Concluding remarks

In the present paper we compared the cost complexities of the approximative truncated singular value decomposition and the general sparse grid approach. We have shown that the sparse grid provides an efficient tool to approximate two-variate functions. Its cost complexity is at least equal to the approximative truncated singular value decomposition. In many situations it is even superior.

In case of the sparse grid approach we envision further improvements by the use of local adaptivity, which would further increase its performance.

In case of the singular value decomposition the truncation length is determined by the smoothness of the function under consideration and is thus fixed. Therefore, improvements for the truncated singular value decomposition can only be achieved by a more efficient representation of the eigenfunctions.

References

- BALESCU, R. (1997) *Statistical Dynamics, Matter out of Equilibrium*. Imperial College Press, Imperial College, London.
- BEBENDORF, M. (2011) Adaptive cross approximation of multivariate functions. *Constr. Approx.*, **34**, 149–179.
- BARRETT, J.W. & KNEZEVIC, D. & SÜLI, E. (2009) Kinetic models of dilute polymers: analysis, approximation and computation. 11th School on Mathematical Theory in Fluid Mechanics 22–29 May 2009, Kacov, Czech Republic Necas Center for Mathematical Modeling, Prague.
- BUNGARTZ, H.J. & GRIEBEL, M. (2004) Sparse grids. *Acta Numerica*, **13**, 147–269.

- CIORANESCU, D. & DAMLAMIAN, A. & GRISO, G. (2008) The periodic unfolding method in homogenization. *SIAM J. Appl. Math.*, **40**, 1585–1620.
- DAHMEN, W. & ROHWEDDER, T. & SCHNEIDER, R. & ZEISER, A. (2008) Adaptive eigenvalue computation: complexity estimates. *Numer. Math.*, **110**, 277–312.
- DAI, X. & XU, J. & ZHOU, A. (2008) Convergence and optimal complexity of adaptive finite element eigenvalue computations. *Numer. Math.*, **110**, 313–355.
- DEVORE, R. & KONYAGIN, S. & TEMLYAKOV, V. (1998) Hyperbolic wavelet approximation. *Constr. Approx.*, **14**, 1–26.
- DYAKONOV, E. (1996) *Optimization in Solving Elliptic Problems*. CRC Press, Boca Raton.
- GHANEM, R.G. & SPANOS, P.D. (1991) *Stochastic Finite Elements: A Spectral Approach*. Springer-Verlag.
- GRASEDYCK, L. (2010) Hierarchical singular value decomposition of tensors. *SIAM J. Matrix Anal. Appl.*, **31**, 2029–2054.
- GRIEBEL, M. (1994) *Multilevelmethoden als Iterationsverfahren über Erzeugendensystemen*. Teubner Skripten zur Numerik. B.G. Teubner, Stuttgart.
- GRIEBEL, M. & HARBRECHT, H. (2011). On the construction of sparse tensor product spaces. Preprint 497, Berichtreihe des SFB 611, Universität Bonn, Germany. (to appear in *Math. Comput.*)
- GRIEBEL, M. & KNAPEK, S. (2009). Optimized general sparse grid approximation spaces for operator equations. *Math. Comput.*, **78**, 2223–2257.
- GRIEBEL, M. & OELTZ, D. (2007). A sparse grid space-time discretization scheme for parabolic problems. *Computing*, **81**, 1–34.
- GRIEBEL, M. & OSWALD, P. (1994). On additive Schwarz preconditioners for sparse grid discretizations. *Numer. Math.*, **66**, 449–463.
- GRIEBEL, M. & OSWALD, P. & SCHIEKOFER, T. (1999). Sparse grids for boundary integral equations. *Numer. Math.*, **83**, 279–312.
- HACKBUSCH, W. (2012). *Tensor spaces and numerical tensor calculus*. Springer, Berlin-Heidelberg.
- HACKBUSCH, W. & KÜHN, S. (2009). A new scheme for the tensor representation. *J. Fourier Anal. Appl.*, **15**(5), 706–722.
- HARBRECHT, H. (2010) A finite element method for elliptic problems with stochastic input data. *Appl. Numer. Math.*, **60**, 227–244.
- HARBRECHT, H. & SCHNEIDER, R. & SCHWAB, C. (2008) Sparse second moment analysis for elliptic problems in stochastic domains. *Numer. Math.*, **109**, 167–188.
- HARBRECHT, H. & SCHNEIDER, R. & SCHWAB, C. (2008) Multilevel frames for sparse tensor product spaces. *Numer. Math.*, **110**, 199–220.

- HOANG, V.H. & SCHWAB, C. (2005) High-dimensional finite elements for elliptic problems with multiple scales. *SIAM Multiscale Model. Simul.*, **3**, 168–194.
- KOLMOGOROV, A. (1936) Über die beste Annäherung von Funktionen einer gegebenen Funktionenklasse. *Annals Math.*, **37**, 107–110.
- LE BRIS, C. & LELIÈVRE, T. (2009) Multiscale modelling of complex fluids: A mathematical initiation. In *Multiscale Modeling and Simulation in Science*, Lecture Notes in Computational Science and Engineering, vol. 66, Springer, Berlin, pages 49–138.
- LÒÈVE, M. (1978) *Probability Theory*, Vol. I+II, Springer, New York.
- OSELEDTS, I.V. & TYRTYSHNIKOV, E.E. (2009) Breaking the curse of dimensionality, or how to use SVD in many dimensions. *SIAM J. Sci. Comput.*, **31**(5), 3744–3759.
- LOZINSKI, A. & OWENS, R.G. & PHILLIPS, T.N. (2010) The Langevin and Fokker-Planck equations in polymer rheology. In R. Glowinski (ed.) *Handbook of Numerical Analysis XVI/XVII*, Elsevier North-Holland.
- MATACHE, A. (2002) Sparse two-scale FEM for homogenization problems. *SIAM J. Sci. Comput.*, **17**, 709–720.
- SCHWAB, C. & TODOR, R.A. (2003) Sparse finite elements for elliptic problems with stochastic loading. *Numer. Math.*, **95**, 707–734.
- SCHWAB, C. & TODOR, R.A. (2003) Sparse finite elements for stochastic elliptic problems. Higher order moments. *Computing*, **71**, 43–63.
- SCHWAB, C. & TODOR, R.A. (2006) Karhunen-Loève approximation of random fields by generalized fast multipole methods. *J. Comput. Phys.*, **217**, 100–122.
- STEVENSON, R. & SCHWAB, C. (2009) Space-time adaptive wavelet methods for parabolic evolution problems. *Math. Comput.*, **78**, 1293–1318.
- STRÖMBERG, J. (1998) Computation with wavelets in higher dimensions. In *Proceedings of the International Congress of Mathematicians, Vol. III, Berlin, 1998*, Doc. Math., Extra Vol. III, 523–532.
- WIDMER, G. & HIPTMAIR, R. & SCHWAB, C. (2008) Sparse adaptive finite elements for radiative transfer. *J. Comp. Phys.*, **227**(12), 6071–6105.
- ZENGER, C. (1991) Sparse grids. In *Parallel algorithms for partial differential equations*, Proceedings of the 6th GAMM-Seminar, Kiel/Germany, 1990, Notes Numer. Fluid Mech. 31, pages 241–251, Vieweg, Braunschweig.